

# GREEN'S FUNCTION MONTE CARLO CALCULATIONS OF LIGHT NUCLEI

S. C. Pieper and R. B. Wiringa

WORK WITH

Joe Carlson, Los Alamos

Vijay Pandharipande, Urbana

K. Varga, Oak Ridge

Work not possible without

Math & Computer Science Division Computers

(Est. 900K CPU-hours  $\approx$  110 TFLOP hours in FY01)

NERSC IBM SP

(963K charge hours = 385K CPU-hours

$\approx$  135 TFLOP hours in FY01)

## Two Problems in Microscopic Few- & Many-Nucleon Calculations

(I) What is the Hamiltonian?

- NN force is reasonably controlled
- 3N force must be determined while computing properties of light nuclei!

(II) Given  $\mathcal{H}$ , solve the Schrödinger equation for  $A$  nucleons accurately.

- Much recent progress for  $A \leq 10$

Direct comparison of calculations to data is ambiguous if (II) is not solved.

Our goal is a microscopic description of nuclear structure and reactions from bare NN & 3N forces and consistent currents.

## NUCLEAR HAMILTONIAN

$$\mathcal{H} = \sum_i K_i + \sum_{i < j} v_{ij} + \sum_{i < j < k} V_{ijk}$$

$v_{ij}$ : Argonne  $v_{18}$

$$v_{ij} = v_{ij}^\gamma + v_{ij}^{CS} + v_{ij}^{CD} \quad v_{ij}^{CS} = \sum_{p=1,14} v_p(r_{ij}) O_{ij}^p$$

$$O_{ij}^{p=1,14} =$$

$$[1, \sigma_i \cdot \sigma_j, S_{ij}, \mathbf{L} \cdot \mathbf{S}, \mathbf{L}^2, \mathbf{L}^2 \sigma_i \cdot \sigma_j, (\mathbf{L} \cdot \mathbf{S})^2] \otimes [1, \tau_i \cdot \tau_j]$$

$V_{ijk}$ : Urbana IX and new Illinois models

Need to solve

$$\mathcal{H}\Psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_A; s_1, s_2, \dots, s_a; t_1, t_2, \dots, t_A)$$

$$= E\Psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_A; s_1, s_2, \dots, s_a; t_1, t_2, \dots, t_A)$$

$s_i$  are nucleon spins:  $\pm \frac{1}{2}$

$t_i$  are nucleon isospins (proton or neutron):  $\pm \frac{1}{2}$

$2^A \times \left( \frac{A}{Z} \right)$  coupled equations in  $3A$  variables

(number of isospin states can be somewhat limited)

## VARIATIONAL MONTE CARLO

Minimize expectation value of  $\mathcal{H}$

$$E_T = \frac{\langle \Psi_T | \mathcal{H} | \Psi_T \rangle}{\langle \Psi_T | \Psi_T \rangle} \geq E_0$$

Simplified trial wave function:

$$|\Psi_T\rangle = [1 + \sum_{i < j < k} U_{ijk}] [\mathcal{S} \prod_{i < j} (1 + U_{ij})] \prod_{i < j} f_{ij} |\Phi\rangle$$

$U_{ijk}$  are 3-body correlations from  $V_{ijk}$

$U_{ij}$  are non-commuting 2-body correlations from  $v_{ij}$

$f_{ij}$  are central (mostly short-ranged repulsion) correlations

$\Phi$  is a  $1\hbar\omega$  shell-model w.f.

- determines quantum numbers of state
- fully antisymmetric
- translationally invariant
- has multiple spatial-symmetry components

## GREEN'S FUNCTION MONTE CARLO

VMC  $\Psi_T$  propagated to imaginary time  $\tau$ :

$$\Psi(\tau) = e^{-(\mathcal{H}-E_0)\tau} \Psi_T$$

$$\Psi_0 = \lim_{\tau \rightarrow \infty} \Psi(\tau)$$

$$\mathcal{H}\Psi_0 = E_0\Psi_0$$

Small time-step propagator:

$$\Psi(\tau) = \left[ e^{-(\mathcal{H}-E_0)\Delta\tau} \right]^n \Psi_T; \quad \tau = n\Delta\tau$$

$$G_{\beta\alpha}(\mathbf{R}', \mathbf{R}) = \langle \mathbf{R}', \beta | e^{-(\mathcal{H}-E_0)\Delta\tau} | \mathbf{R}, \alpha \rangle$$

$$\Psi(\mathbf{R}_n, \tau) =$$

$$\int G(\mathbf{R}_n, \mathbf{R}_{n-1}) \cdots G(\mathbf{R}_1, \mathbf{R}_0) \Psi_T(\mathbf{R}_0, 0) d\mathbf{R}_{n-1} \cdots d\mathbf{R}_0$$

Fermion sign problem limits maximum  $\tau$ :

$G$  brings in lower-energy boson solution

$\langle \Psi_T | \mathcal{H} | \Psi(\tau) \rangle$  projects back fermion solution

Exponentially growing statistical errors

Constrained-path propagation, suggested by J. Carlson,  
removes steps that have

$$\overline{\Psi(\tau, \mathbf{R})^\dagger \Psi(\mathbf{R})} = 0$$

Many tests demonstrate reliability

## MAKING IT PARALLEL

Master-slave structure

Each slave gets configurations to propagate

Results sent back to master for averaging as generated

During propagation, configs multiply or are killed

- Work load fluctuates
- Periodically master collects statistics and tells slaves to redistribute
- Slaves have work set aside to do during this synchronization

Large calculations have very low (minutes) frequency of communication

Parallelization efficiencies typically 95%

92% efficiency obtained on 2048-processor Seaborg run; 0.55 TFLOPS.

## TYPICAL CURRENT CALCULATIONS

- Propagation to  $\tau = 0.2 - 0.4 \text{ MeV}^{-1}$
- $E(\tau)$  every  $\tau = 0.01 \text{ MeV}^{-1}$  (0.02 for  $A \geq 10$ )
- Average of  $E(\tau)$  for  $\tau \geq 0.1$

	Config- urations	$\tau_{\text{max}}$ $\text{MeV}^{-1}$	Statistical Error (MeV)	Processor hours*
${}^6\text{Li}$	50,000	0.2	0.08	40
${}^7\text{Li}$	20,000	0.2	0.14	340
${}^8\text{Be}$	15,000	0.2	0.2	300
${}^8\text{Li}$	12,000	0.2	0.2	600
${}^9\text{Be}$	6,500	0.4	0.5	10,000
${}^9\text{Li}$	8,000	0.4	0.4	13,500
${}^{10}\text{B}$	5,000	0.5	0.5	5,000
${}^{10}\text{Be}$	3,000	0.6	0.6	9,000

- \*6 – 8: IBM SP3 or SGI 250 MHz R10000 processors  
 9: 500 MHz Pentium-III at  $\sim 110$  MFLOPS  
 (MCS Chiba City)  
 10: IBM SP at  $\sim 320$  MFLOPS (NERSC Seaborg)

# ${}^3\text{H}(\alpha, \gamma){}^7\text{Li}$ & ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$ CAPTURE REACTIONS

U. of Chicago thesis work of K. Nollett

Source of  ${}^7\text{Li}$  in the big bang

- Astrophysically important region is 20–500 keV.


${}^7\text{Be}$  reaction also source of solar neutrinos

- Astrophysically important region is 20 keV.
- No data in this region

Full 7-nucleon calculation

- $A = 7$  wave functions have proper 3+4 cluster form.

${}^2\text{H}(\alpha, \gamma){}^6\text{Li}$  also done





## RECENT PROGRESS

- New computers and methods allow  $\sim 1 - 2\%$  calculations of light p-shell nuclear energies
- First  $A = 9, 10$  calculations done in FY2000. Full set done in 2001 to early 2002
- First  $A = 10$  unnatural-parity states done in FY2001
- Extensions to the Urbana  $V_{ijk}$  give average binding-energy errors  $< 0.7$  MeV for  $A = 3 - 10$  nuclei
- Variational w.f. with proper asymptotic clustering
- VMC calculations of weak transitions in  $A = 6, 7$
- Participated in  $^4\text{He}$  benchmark paper using seven precise methods; GFMC good to  $\leq 0.1\%$  for AV8'
- Study of what is needed in nuclear potentials to get observed nuclear level structure

## PLANS

- GFMC for scattering states – widths of resonances
- $A = 9, 10$  – continue study of intruder states
- ${}^7\text{Be}(p, \gamma){}^8\text{B}$ ,  ${}^8\text{B}(\beta^+){}^8\text{Be}$ ,  ${}^9\text{Be}(e, e'\text{N})$ ,  ${}^{10}\text{C}(\beta^+){}^{10}\text{B}$ , ...
- neutron-rich systems – input for large-nuclei methods
- ${}^{12}\text{C}$  by GFMC (several years away)

We are approaching a nuclear standard model for computing nuclear properties and reactions

GFMC calculations are the benchmark for  $6 \leq A \leq 10$

